

Evidence for a positive cosmological constant from flows of galaxies and distant supernovae

Idit Zehavi^{*,†} & Avishai Dekel^{*}

^{*}Racah Institute for Physics, The Hebrew University, Jerusalem 91904, Israel

[†]NASA/Fermilab Astrophysics Group, Fermi National Accelerator Laboratory, Box 500, Batavia, IL 60510-0500, USA

Recent observations^{1,2} of high-redshift supernovae seem to suggest that the global geometry of the Universe may be affected by a ‘cosmological constant’, which acts to accelerate the expansion rate with time. But these data by themselves still permit an open universe of low mass density and no cosmological constant. Here we derive an independent constraint on the lower bound to the mass density, based on deviations of galaxy velocities from a smooth universal expansion^{3–7}. This constraint rules out a low-density open universe with a vanishing cosmological constant, and together the two favour a nearly flat universe in which the contributions from mass density and the cosmological constant are comparable. This type of universe, however, seems to require a degree of fine tuning of the initial conditions that is in apparent conflict with ‘common wisdom’.

The standard cosmological model based on Einstein’s gravity and global homogeneity is characterized by two fundamental parameters that measure the main contributions to the total energy density (in terms of a ‘critical’ density): the mean mass density Ω_m and the cosmological constant Ω_Λ . The latter, an intrinsic part of the theory of general relativity, represents a uniform energy density that is associated with the vacuum (not the mass), and, if positive, acts as a repulsive component of the global gravitational force field. The values of these parameters determine the type of the universe we live in, its global extent and ultimate fate, as follows. The parameter plane (Fig. 1) is divided by three lines into six regions. The line $\Omega_m + \Omega_\Lambda = 1$ defines a ‘flat’ euclidean geometry, favoured by theories of inflation in the early universe; it separates curved-space (non-euclidean) models of ‘closed’ finite extent (above) and ‘open’ infinite extent (below). The horizontal line $\Omega_\Lambda = 0$ roughly distinguishes between an ‘unbound’ universe that will expand forever (above) and a ‘bound’ universe that will eventually re-collapse (below). The special point $(\Omega_m, \Omega_\Lambda) = (1, 0)$ corresponds to the simplest model termed Einstein-deSitter; unless the universe exactly fits this model, it

evolves away from this point as it expands. If $\Omega_\Lambda > 0$, the universe eventually crosses the line $\Omega_\Lambda = 0.5\Omega_m$, which separates between a universe whose expansion decelerates by attraction (below) or accelerates by repulsion (above).

New techniques to measure distances of supernovae (type Ia) at large distances (cosmological redshifts $0.5 - 1$), where the large-scale curvature of space-time plays a noticeable role, enable a classical cosmological test based on how the time dependence of the Hubble relation between velocity and distance depends on the cosmological parameters⁸. The corresponding, inclined (blue) confidence limits in Fig. 1 are based on the results of the Supernova Cosmology Project¹, which are fully consistent with the findings of the High- z Supernova Search Team². The allowed region in the parameter plane is an elongated stripe, crudely approximated by $0.8\Omega_m - 0.6\Omega_\Lambda = -0.2 \pm 0.1$ (1σ errors). Based on this result alone, a nearly flat universe is likely, with a broad maximum to the probability near $(\Omega_m, \Omega_\Lambda) = (0.3, 0.7)$, but an open model, of $(0.1, 0)$, for example, is still allowed at the 2% confidence level. An orthogonal constraint is required in order to remove the degeneracy between Ω_Λ and Ω_m .

Here we show that such a constraint is provided by current data of ‘peculiar’ velocities of galaxies on scales of ~ 300 million light years. These velocities, which are produced by gravity due to local fluctuations about the mean mass density, depend also on the value of the mean density itself, Ω_m . The constraints on Ω_m from these velocities are almost independent of Ω_Λ , but combined with the inclined supernovae constraints any bound on Ω_m effectively serves as a bound on Ω_Λ .

We use two recent catalogues of galaxy peculiar velocities. One, named Mark III⁹, contains distances for $\sim 3,000$ galaxies within $\sim 70 h^{-1}\text{Mpc}$; the other, named SFI^{10,11}, consists of $\sim 1,300$ spiral galaxies with a more uniform sampling in a similar volume. In order to obtain the peculiar velocity of a galaxy, its total velocity is measured using the Doppler redshift, and its distance is separately inferred using the so-called Tully-Fisher method, with an error of 15 – 21%. The desired peculiar velocity along the line of sight is obtained by subtracting, from the total velocity, the Hubble expansion velocity at the galaxy distance. The peculiar velocities are carefully corrected for systematic errors^{7,12}.

The POTENT method^{12,13} assumes that the peculiar velocities were generated by gravity and that they therefore represent a potential flow on large scales. This allows a recovery of the three-dimensional velocity field of galaxies in our local cosmological neighbourhood. For an assumed value of Ω_m , the underlying field of mass-density fluctuations can be extracted, using slightly nonlinear approximations to the relation between velocity and density fluctuations within the theory of gravitational instability. Assuming also that the initial fluctuations were a gaussian random field, direct dynamical constraints on Ω_m can be obtained without appealing to the observed spatial distribution of galaxies. These constraints are thus independent of the unknown relation between galaxies and the underlying mass density (termed ‘biasing’). Several different methods applied to the POTENT Mark III fields consistently yield a lower bound of $\Omega_m > 0.3$ at the 99% confidence level, marked by a vertical red line in Fig. 1. One method constrains Ω_m based on the large diverging outflow




Figure 1: Constraints in the $\Omega_m - \Omega_\Lambda$ plane, showing relative confidence limits of 68%, 90% and 99%. The inclined contours (blue, same in both panels), that roughly constrain $0.8\Omega_m - 0.6\Omega_\Lambda$, arise from the global geometry of space-time based on supernovae as distance indicators¹. The lower bound $\Omega_m > 0.3$ (red) represents a 99% confidence limit based on several different studies of peculiar velocities using POTENT reconstruction, independent of the biasing relation between galaxies and mass³⁻⁵. The almost vertical contours (red) arise from the peculiar velocities of the Mark III (left) and SFI (right) catalogues, based on a likelihood analysis which assumes a parametric mass power spectrum of the CDM family with COBE normalization on large scales^{6,7}. The central contours shown are for a standard case of fixed $h = 0.65$ and $n = 1$; the uncertainties in these values are represented by the outer 99% contours on both sides (long dashed red), computed for the extreme cases of fixed $(h, n) = (0.55, 0.9)$ and $(0.75, 1.1)$. These encompass a conservative range of non-negligible likelihood based on the flow data. The strong bounds from flows are on Ω_m ; the apparent upper bound on Ω_Λ arises indirectly from the COBE normalization and is very uncertain; the contour can extend further in the Ω_Λ direction, allowing significant overlap with the supernova confidence region. The corresponding joint (relative) confidence limits from supernovae and flows are shown (black contours); they are almost the same for Mark III and SFI. The combined constraints thus favour an unbound and roughly flat universe with comparable contributions from the cosmological constant and mass density. A universe with very low Ω_m and zero cosmological constant is ruled out.

recovered in the vicinity of a large nearby underdense region (void), using the fact that even an empty void cannot induce outflows as large as those observed if the density outside it is too low³. A second method uses the deviations of the initial fluctuations from a gaussian distribution, when derived from the velocity data, assuming a value of Ω_m that is too low⁴. A third method derives Ω_m based on the gravitationally-induced deviations from gaussian distribution of the present-day velocity-divergence field⁵. Similar constraints confirming the lower bound of $\Omega_m > 0.3$ are currently being obtained from the SFI data.

In a new analysis^{6,7}, not based on POTENT, we have applied a maximum-likelihood analysis to the Mark III and SFI peculiar velocities to determine cosmological parameters. It is done via a parametric model for the mass-density fluctuation power spectrum P_k , which measures the distribution of power among different length scales, proportional to k^{-1} . This analysis uses linear gravitational theory and assumes that both the fluctuations and the errors are gaussian. The model used is of the popular cold dark matter (CDM) model family; P_k is normalized¹⁴ by the fluctuations in the cosmic microwave background (CMB) as measured on very large scales by the COBE satellite. The result provides constraints on permissible combinations of three parameters: Ω_m , the power index n (where $P_k \propto k^n$ on large scales), and the Hubble expansion constant h . These parameters enter via the shape of P_k as well as from the geometry and dynamics of space-time. The constraints obtained from the two data sets turn out to be similar; they define a two-dimensional surface in the Ω_m - h - n space. In the case of a flat cosmology this surface can be crudely approximated by $\Omega_m h_{65}^{1.3} n^2 = 0.58 \pm 0.12$ (where h_{65} is the Hubble constant in units of $65 \text{ km s}^{-1} \text{ Mpc}^{-1}$, and the errors refer to 90% uncertainty).

The obtained P_k is determined by the velocity data (on scales of $\sim 50 \text{ h}^{-1} \text{ Mpc}$) and is not much affected by the COBE normalization (at $\sim 1,000 \text{ h}^{-1} \text{ Mpc}$); a similar P_k is reproduced to within a few percent when the amplitude on large scales is left as a free parameter. Also, the result is not dominated by the assumed distance errors; P_k is reproduced to better than 10% when an error model with free parameters is incorporated in the likelihood analysis⁷. This robustness is within the 1σ limits of the likelihood analysis.

The peculiar velocities have also been analysed by methods that incorporate the spatial distribution of galaxies. For example, a comparison of the Mark III velocities and the IRAS 1.2Jy redshift survey¹⁵ yields results roughly consistent with the CDM model of $\Omega_m \simeq 0.5$ favoured by the P_k analysis of flows reported above. Our current bounds on Ω_m from flows also partly overlap with constraints from the evolution of galaxy clusters¹⁶ ($\Omega_m = 0.45 \pm 0.2$). However, a comprehensive joint analysis of all the available constraints is beyond the scope of this work. In particular, a thorough interpretation of the results involving redshift surveys (which span a noticeable range) must include a full discussion of the non-trivial biasing relation between galaxies and mass¹⁷, and of the complex systematic effects involved in these different studies. Instead, we choose to focus here on one set of constraints, the dynamical constraints from flows, combined with the supernova constraints.

In order to obtain the desired constraints in the Ω_m - Ω_Λ plane, we have applied the P_k

likelihood analysis to the Mark III and SFI velocities, using the COBE-normalized¹⁸ CDM models with varying Ω_m and Ω_Λ that now span the whole parameter plane. The velocities are expected to be insensitive to the value of Ω_Λ ¹⁹, but a weak Ω_Λ dependence occurs as a result of the COBE normalization imposed. This is responsible for the slight left bending of the red vertical likelihood ridges (Fig. 1), and the apparent upper bounds on Ω_Λ . The 1σ error in the COBE normalization translates to additional uncertainties of $\pm 6\%$ in Ω_m and $\pm 20\%$ in Ω_Λ , which are on the order of the uncertainties displayed by the likelihood contours shown. Thus, the COBE errors would tend to stretch the likelihood contours vertically along the Ω_Λ axis, weakening the apparent upper bounds on Ω_Λ .

Since the velocity data favour an extended two-dimensional surface in the Ω_m - h - n space, the likelihood analysis cannot determine the three parameters simultaneously; two of them should be fixed initially. We adopt as our standard case the simplest, scale-invariant ($n = 1$) initial spectrum, and $h = 0.65$ (as favoured by nearby supernova data²⁰). We crudely estimate a $\pm 15\%$ uncertainty in the Hubble constant²¹, and $\pm 10\%$ in the power index²². In order to illustrate the sensitivity of the constraints to these parameters, we have also determined Ω_m and Ω_Λ for two extreme cases of h and n values. The outer 99% contours are shifted accordingly in Fig. 1.

The likelihood ridge of SFI extends into higher values of Ω_Λ than for Mark III. The independent constraints from flows (using P_k) and supernovae overlap significantly for SFI, but only near the 90% contours for Mark III with $(h, n) = (0.65, 1)$ (which improves when COBE errors are considered or if $n > 1$ or $h > 0.65$). The large uncertainty in the upper bound on Ω_Λ indicates that the degree of overlap should not be interpreted as indicating either consistency or inconsistency between the velocity and supernova data and the assumed standard cosmological model.

A joint parameter estimation from supernovae and flows is therefore meaningful, although it is limited to relative likelihoods; a measure of absolute probabilities is not straightforward, despite a seemingly acceptable goodness of fit. The black joint contours (Fig. 1) are computed by multiplying the two likelihood values at each point, having assumed that the data are independent. The most likely joint values (shown as crosses in Fig. 1) for the supernovae and the combined Mark III and SFI dataset are $(\Omega_m, \Omega_\Lambda) = (0.5, 0.8)$, while the conservative joint 99% confidence limits allow Ω_m values in the range $0.3 - 0.9$ and Ω_Λ values in the range $0.1 - 1.4$. The constraints obtained separately from Mark III and SFI are very similar.

The constraints from local flows thus remove the degeneracy in the constraints from the global-geometry test based on supernovae (and vice versa), and help rule out an open model with zero cosmological constant. A nearly flat universe, with comparable contributions of matter and cosmological constant to the total energy density, is likely. The favoured model is therefore an unbound universe that will accelerate forever, although it is impossible yet to determine whether the global geometry is flat, open or closed.

Such comparable contributions from the mass density and the cosmological constant represent a puzzling fine tuning, for example, because the two parameters, Ω_m and Ω_Λ , are

expected to vary with time in opposite senses. The standard theory expects the cosmological constant either to vanish or be larger by many orders of magnitude^{8,23}. Perhaps entropic arguments may be needed to explain such a fine tuning. Although the observed constraints are not yet finally confirmed, they already seem interesting enough to pose a serious challenge to theoretical physics.

Other constraints in the Ω_m - Ω_Λ plane are worth mentioning in this framework. Constraints consistent with the supernova ridge but of larger uncertainty arise from the age of old star clusters²⁴ versus the Hubble expansion rate. Constraints of orthogonal orientation, roughly on $\Omega_\Lambda + \Omega_m$, can be deduced from the acoustic peaks in the sub-degree angular power spectrum of fluctuations in the CMB, as observed from balloons and from the ground^{22,25–29}. Two CMB satellites planned for the next decade, MAP and Planck, are expected to provide more accurate constraints^{27,30}.

Future peculiar-velocity data are expected to improve the accuracy of the constraints. In addition to many new velocities based on distance indicators of the Tully-Fisher type, the most promising sources of large-scale peculiar velocities in the future will probably be the siblings of the same supernovae discussed above, but at low redshifts; their distances can be measured with 5 – 10% accuracy out to large distances and their sampling density is limited only by the patience of the observers.

References

- [1] Perlmutter, S. *et al.* Measurements of Ω and Λ from 42 high-redshift supernovae. *Astrophys. J.* **517**, 565-586 (1999).
- [2] Riess, A.G. *et al.* Observational evidence from supernovae for an accelerating universe and a cosmological constant. *Astron. J.* **116**, 1009-1038 (1998).
- [3] Dekel, A. & Rees, M.J. Ω from velocities in voids. *Astrophys. J.* **422**, L1-L4 (1994).
- [4] Nusser, A. & Dekel, A. Ω and the initial fluctuations from velocity and density fields. *Astrophys. J.* **405**, 437-448 (1993).
- [5] Bernardeau, F., Juszkiewicz, R., Dekel, A. & Bouchet, F.R. Omega from the skewness of the cosmic velocity divergence. *Mon. Not. R. Astron. Soc.* **274**, 20-26 (1995).
- [6] Zaroubi, S., Zehavi, I., Dekel, A., Hoffman, Y. & Kolatt T. Large-scale power spectrum from peculiar velocities via likelihood analysis. *Astrophys. J.* **486**, 21-31 (1997).
- [7] Freudling, W., Zehavi, I., da Costa, L.N., Dekel, A., Eldar, A., Giovanelli, R., Haynes, M.P., Salzer, J.J., Wegner, G. & Zaroubi, S. Large-scale power spectrum and cosmological parameters from SFI peculiar velocities. *Astrophys. J.* **523**, 1-15 (1999).
- [8] Carroll, S.M., Press, W.H. & Turner, E.L. The cosmological constant. *Annu. Rev. Astron. Astrophys.* **30**, 499-542 (1992).

- [9] Willick, J.A., Courteau, S., Faber, S.M., Burstein, D., Dekel, A. & Strauss, M.A. Homogeneous velocity-distance data for peculiar velocity analysis. III. The Mark III catalog of galaxy peculiar velocities. *Astrophys. J. S.* **109**, 333-366 (1997).
- [10] Haynes, M.P., Giovanelli, R., Salzer, J.J., Wegner, G., Freudling, W., da Costa, L.N., Herter, T. & Vogt, N.P. The I-band Tully-Fisher relation for Sc galaxies: optical imaging data. *Astron. J.* **117**, 1668-1687 (1999).
- [11] Haynes, M.P., Giovanelli, R., Chamaraux, P., da Costa, L.N., Freudling, W., Salzer, J.J. & Wegner, G. The I-band Tully-Fisher relation for Sc galaxies: 21 centimeter HI line data. *Astron. J.* **117**, 2039-2051 (1999).
- [12] Dekel, A., Eldar, A., Kolatt, T., Willick, J.A., Faber, S.M., Courteau, S. & Burstein, D. POTENT reconstruction from Mark III velocities. *Astrophys. J.* **522**, 1-38 (1999).
- [13] Dekel, A., Bertchinger, E. & Faber, S.M. Potential, velocity, and density fields from sparse and noisy redshift-distance samples: Method. *Astrophys. J.* **364**, 349-369 (1990).
- [14] Sugiyama, N. Cosmic background anisotropies in cold dark matter cosmology. *Astrophys. J. S.* **100**, 281-305 (1995).
- [15] Sigad, Y., Eldar, A., Dekel, A., Strauss, M.A. & Yahil, A. IRAS versus POTENT Density Fields on Large Scales: Biasing and Omega. *Astrophys. J.* **495**, 516-532 (1998).
- [16] Eke, V.R., Cole, S., Frenk, C.S. & Henry, P.J. Measuring Ω_0 using cluster evolution. *Mon. Not. R. Astron. Soc.* **298**, 1145-1162 (1998).
- [17] Dekel, A. & Lahav, O. Stochastic nonlinear galaxy biasing. *Astrophys. J.* **520**, 24-34 (1999).
- [18] Bunn, E.F. & White, M. The four-year COBE normalization and large-scale structure. *Astrophys. J.* **480**, 6-21 (1997).
- [19] Lahav, O., Lilje, P.B., Primack, J.R. & Rees, M.J. Dynamical effects of the cosmological constant. *Mon. Not. R. Astron. Soc.* **251**, 128-136 (1991).
- [20] Riess, A.G., Press, W.H. & Kirshner, R.P. A precise distance indicator: type Ia supernova multicolor light-curve shape. *Astrophys. J.* **473**, 88-109 (1996).
- [21] Freedman, W.L. Determination of the Hubble constant. in *Critical Dialogues in Cosmology* (ed Turok, N.) 92-129 (Singapore, World Scientific, 1997).
- [22] Bond, J.R. & Jaffe, A.H. Constraining large scale structure theories with the cosmic background radiation. *Phil. Trans. R. Soc. Lond. A* (in the press); also as preprint astro-ph/9809043 (1998).
- [23] Weinberg, S. The cosmological constant problem. *Rev. Mod. Phys.* **61**, 1-23 (1989).
- [24] Chaboyer, B., Demarque, P. Kernan, P.J. & Krauss, L.M. The age of globular clusters in light of Hipparcos: Resolving the age problem? *Astrophys. J.* **494**, 96-110 (1998).

- [25] Efstathiou, G., Bridle, S.L., Lasenby, A.N., Hobson, M.P. & Ellis, R.S., Constraints on Ω_Λ and Ω_m from distant type Ia supernovae and cosmic microwave background anisotropies. *Mon. Not. R. Astron. Soc.* **303**, L47-L52 (1999).
- [26] Lineweaver, C.H. The cosmic microwave background and observational convergence in the $\Omega_m - \Omega_\Lambda$ plane. *Astrophys. J.* **505**, L69-L73 (1998).
- [27] Tegmark, M., Eisenstein, D.J., Hu, W. & Kron, R. Cosmic complementarity: probing the acceleration of the universe. *Astrophys. J.* submitted; also as preprint astro-ph/9805117 (1998).
- [28] White, M. Complementary measures of the mass density and cosmological constant. *Astrophys. J.* **506**, 495-501 (1998).
- [29] Lasenby, A.W., Jones, A.W. & Dabrowski, Y. Observations of the cosmic microwave background and implications for cosmology and large scale structure. *Phil. Trans. R. Soc. Lond. A* (in the press); also as preprint astro-ph/9810196 (1998).
- [30] Zaldarriaga, M., Spergel, D.N. & Seljak, U. Microwave background constraints on cosmological parameters. *Astrophys. J.* **488**, 1-13 (1997).

Acknowledgements. We thank the Mark III and SFI teams, including D. Burstein, S. Courteau, L.N. daCosta, A. Eldar, S.M. Faber, W. Freudling, R. Giovanelli, M.P. Haynes, T. Kolatt, J.J. Salzer, M.A. Strauss, G. Wagner, J.A. Willick, A. Yahil, & S. Zaroubi. We thank G.R. Blumenthal and O. Lahav for discussions. This research was supported by US-Israel Binational Science Foundation grant 95-00330 and Israel Science Foundation grant 546/98 at the Hebrew University, and by the DOE and the NASA grant NAG 5-7092 at Fermilab.

Correspondence and requests for materials should be addressed to I.Z. (email: iditz@fnal.gov) and A.D. (email: dekel@astro.huji.ac.il).